arXiv:cond-mat/9810267v1 [cond-mat.mes-hall] 21 Oct 1998

Edge Occupation Numbers of a Half-Filled Landau Level

J. H. Han[1] and S.-R. Eric Yang[1,2]

Asia Pacific Center for Theoretical Physics, 207-43 Cheongryangri-dong Dongdaemun-gu Seoul 130-012, Korea[1]

and

Department of Physics, Korea University, Seoul 136-701, Korea[2]

We demonstrate that a theory of the edge of a half-filled Landau level recently proposed by Lee and Wen predicts results for the edge occupation number similar to those of a variational trial wave function proposed previously by us. We treat Lee and Wen's edge action of a half-filled Landau level within the framework of bosonization theory, and show that the momentum occupation numbers are determined by a product of two Green's functions, one charged and one neutral. In the bulk region ($k < 0$) we find a linear occupation profile, $n_k \propto A + Bk$, while in the tail region ($k > 0$) it is exponentially decaying over the range $k \sim \Lambda_n$, the momentum cutoff for neutral mode. We find a good fit with the numerical results for occupation numbers.

PACS numbers: 73.20.Dx, 73.40.Hm

Recently there has been a great deal of interests in two-dimensional electron liquids in a strong magnetic field at half-filling, and several theories have been proposed[1–7]. Unlike insulating QH liquids this new state is metallic. Although all these theories describe a metallic state it is not clear whether they predict identical edge properties. Field theoretical methods of Shytov *et al.*[8] and Lee and Wen[9] seem to predict different dynamic properties of the edge. It is also not clear whether the edge properties of a variational trial wave function of this liquid[5] differs from those predicted by field theories.

After attaching two flux quanta to each electron to realize the Chern-Simons transformation at half-filling, one is apparently left with a Fermi sea without external magnetic field. One can alternatively regard it as $N/2$ of the composite fermions seeing an upward $N/2$ quanta of flux, while the other half sees magnetic flux in the opposite direction, in an N-particle system. On average the net residual flux will be zero. As a consequence half of the composite fermions in the system occupy the "lowest Landau level" (as judged by the composite fermions), and the other half the anti-holomorphic lowest Landau level. This consideration naturally leads to a trial wave function at half-filling in particular, and wave functions at other hierarchy filling factors in general[10,5].

Once a trial wave function for the half-filled Landau level[5] is known the edge occupation number can be computed in the same manner as the calculation of the occupation number from Laughlin's wave function[11]. It turned out that the edge of a metallic liquid has a rather different property than that of incompressible QH states at filling factor $\nu = 1/M$ (M odd integer). It is well known that for incompressible QH states at $\nu = 1/M$ the electron occupation number n_k near the edge is $n_k \propto (k_{ed} - k)^{M-1}\theta(k_{ed} - k)$, where k_{ed} is the wavenumber corresponding to the location of the edge[12,11]. Our result for the quantum Hall droplet at half filling showed that a tail exists in the occupation number beyond the radius of a droplet of uniform filling factor 1/2, i.e. $n_k \neq 0$ for $k - k_{ed} > 0$. This apparent increase in the size of the droplet is due to the displaced zeroes of wave function from the electrons[10]. The occupation numbers were found to exhibit Friedel type oscillations, consistent with the metallic nature of the liquid[10].

Recently Lee and Wen have proposed a two-mode model for the edge[9] based on a theory of the electron liquid at $\nu = 1/2$ consisting of charge and flux liquids[6]. In this theory the charge and the neutral modes of edge deformations are assumed to be well defined up to the momentum cutoffs $\Lambda_{c,n}$, beyond which modes become over-damped. Lee and Wen have calculated tunneling coefficient using this theory, and have found that it agrees with the experimental data of Grayson *et al.*[13]

In this paper we demonstrate that Lee and Wen's theory predicts the edge occupation number profile similar to what is found from the trial wave function[5]. We treat their proposed action within the framework of bosonization theory and find that $n_k \propto A + Bk$ for $-\Lambda_c \ll k < 0$ and $n_k \propto e^{-k/\Lambda_n}$ for $k > 0$ (k is measured from k_{ed}). Our trial wave function also gives linearly increasing occupation numbers for small negative k.

When a proper action is given for the edge at half-filing the electron Green's function can be calclulated. The edge action of a half-filled Landau level derived from the dipole theory of Lee[6] is

$$S_e = \frac{1}{2\pi}\int \frac{d\phi}{dx}\left(\frac{d\phi}{dt} - v_c \frac{d\phi}{dx}\right) dxdt - \frac{1}{4\pi}\int \frac{d\eta}{dx}\left(\frac{d\eta}{dt} + v_n \frac{d\eta}{dx}\right) dxdt + \int (2\phi - \eta)(\vec{j}_t.\hat{n} - \partial_\mu j^\mu) dxdt. \tag{1}$$

It is important to note that the neutral mode is counter-propagating compared with the charged mode, as a result of the minus sign that comes in the $\partial_t\eta\partial_x\eta$ part of the action and the positive-definiteness of the Hamiltonian. The last term in Eq. (1) is the coupling of the edge mode with the transverse bulk current ($\vec{j}_t.\hat{n}$) or the tunneling ($\partial_\mu j^\mu$) current. We parameterize the sample boundary with x, using $\hat{n}$ to indicate the outward normal to the boundary. We quantize the edge action in the absence of the coupling to current sources. The canonical commutation relations for the fields ϕ and η are

$$[\frac{d\phi}{dx}, \phi(y)] = \pi i \delta(x-y); \quad [\frac{d\eta}{dx}, \eta(y)] = -2\pi i \delta(x-y). \tag{2}$$

By comparison with the chiral Luttinger liquid theory at other filling factors[12,14], one may associate $2d\phi/dx$ with the charge displacement at the edge of the liquid. Since the one-dimensional density is $(\nu/2\pi)$(displacement), electron creation/annihilation operators at the boundary will need to satisfy the following algebra:

$$[\frac{\nu}{2\pi}(\text{displacement}), (\text{electron op.})] = \pm\delta(x-y)(\text{electron op.}) \tag{3}$$

One can easily show that $e^{\pm 2i\phi}$ satisfies the desired algebra, but when one interchanges two such operators at x and x', one finds that they obey Bose, rather than Fermi, statistics. To fix the statistics without altering the relation in Eq. (3), we use the following annihilation/creation operators

$$\psi(x) \sim e^{-2i\phi(x) \pm i\eta(x)}, \quad \bar{\psi}(x) \sim e^{2i\phi(x) \mp i\eta(x)}. \tag{4}$$

In Lee and Wen's paper, a specific combination $2\phi - \eta$ appears in the electron Green's function, while our formulation seem to favor both signs. One presumably needs the coupling term in Eq. (1) in order to discriminate the two signs. It will be clear that all the computations below will be insensitive to the choice of sign. We introduce the mode expansion of the boson fields:

$$\begin{aligned} \phi(x) &= \phi_0 + f_0 x + \sum_{n>0} \frac{i}{\sqrt{2n}}[a_n^\dagger e^{-inx} - a_n e^{inx}]; \; [f_0, \phi_0] = i/2, \\ \eta(x) &= \eta_0 + g_0 x + \sum_{n>0} \frac{i}{\sqrt{n}}[b_n e^{-inx} - b_n^\dagger e^{inx}]; \quad [g_0, \eta_0] = -i. \end{aligned} \tag{5}$$

The ground state is defined by the action of the annihilation operators, $a_n|gs\rangle = b_n|gs\rangle = 0$. The time dependence is given by setting $x \to x - v_c t$ for the charged mode, and $x \to x + v_n t$ for the neutral mode.

Since this is a bosonization theory, the phonon Green's functions determine the underlying electron Green's function as well. Define $z \equiv x - v_c t$, and $w \equiv x + v_n t$. We have

$$\begin{aligned} G_\phi(z) &= 4\langle\phi(z)\phi(0)\rangle - 4\langle\phi(0)\phi(0)\rangle = 2\int_0^\infty \frac{dk}{k}[e^{ikz} - 1]e^{-k/\Lambda_c} = -2\ln[1 - i\Lambda_c z], \\ G_\eta(w) &= \langle\eta(w)\eta(0)\rangle - \langle\eta(0)\eta(0)\rangle = -\ln[1 + i\Lambda_n w]. \end{aligned} \tag{6}$$

Electron Green's functions are obtained as

$$\begin{aligned} \langle\bar{\psi}(z)\psi(0)\rangle &= \langle e^{2i\phi(z)} e^{-2i\phi(0)}\rangle\langle e^{-i\eta(z)} e^{i\eta(0)}\rangle = \exp[G_\phi(z) + G_\eta(w)], \\ \langle\psi(0)\bar{\psi}(z)\rangle &= \langle e^{-2i\phi(0)} e^{2i\phi(z)}\rangle\langle e^{i\eta(0)} e^{-i\eta(z)}\rangle = \exp[G_\phi(-z) + G_\eta(-w)]. \end{aligned} \tag{7}$$

It also follows from the definition in Eq. (4) that $\langle\psi(z)\bar{\psi}(0)\rangle = \langle\bar{\psi}(z)\psi(0)\rangle$. Putting our results together,

$$\begin{aligned} \langle\psi(z)\bar{\psi}(0)\rangle &= \langle\bar{\psi}(z)\psi(0)\rangle = 1/[1 - i\Lambda_c z]^2[1 + i\Lambda_n w], \\ \langle\psi(0)\bar{\psi}(z)\rangle &= \langle\bar{\psi}(0)\psi(z)\rangle = 1/[1 + i\Lambda_c z]^2[1 - i\Lambda_n w]. \end{aligned} \tag{8}$$

The time-ordered Green's function $G_e(x,t) = \theta(t)\langle\bar{\psi}(z)\psi(0)\rangle - \theta(-t)\langle\psi(0)\bar{\psi}(z)\rangle$ is given by

$$G_e(x,t) = \frac{\theta(t)}{[1 - i\Lambda_c z]^2[1 + i\Lambda_n w]} - \frac{\theta(-t)}{[1 + i\Lambda_c z]^2[1 - i\Lambda_n w]}. \tag{9}$$

One can consider another kind of Green's function, $G_b(x,t) = \exp[\theta(t)\{G_\phi(z) + G_\eta(w)\} + \theta(-t)\{G_\phi(-z) + G_\eta(-w)\}]$:

$$G_b(x,t) = \frac{\theta(t)}{[1 - i\Lambda_c z]^2[1 + i\Lambda_n w]} + \frac{\theta(-t)}{[1 + i\Lambda_c z]^2[1 - i\Lambda_n w]}. \tag{10}$$

This function is clearly bosonic in character, $G_b(-x,-t) = +G_b(x,t)$. On the other hand, the electron Green's function satisfies $G_e(-x,-t) = -G_e(x,t)$. Notice that one has the approximate symmetry $G_e(x,-t) \approx -G_e(x,t)$ provided $\Lambda_c x, \Lambda_n v_n t \ll 1$, because we have, in such a limit,

$$G_e(x,t) \approx \frac{sgn(t)}{t^2}. \tag{11}$$

We will assume that the momentum cutoffs satisfy $\Lambda_n \ll \Lambda_c$. For the bosonic Green's function $G_b(x,t)$, one observes a similar symmetry in the slightly different limit $\Lambda_n v_n t \ll 1 \ll \Lambda_n x$, and $x \ll v_c t$;

$$G_b(x,t) \approx \frac{1}{t^2}\left(\frac{\theta(t)}{i\Lambda_n x} - \frac{\theta(-t)}{i\Lambda_n x}\right) = \frac{sgn(t)/t^2}{i\Lambda_n x} \approx -G_b(x,-t). \tag{12}$$

The above expression embodies the symmetry that was extensively discussed and emphasized by Lee and Wen[9]. It says that the desired statistics property shows up at distances short compared with $v_c t$, but larger than $1/\Lambda_n$.

We now consider the static property of the edge, in particular its momentum occupation. The momentum distribution is simply a Fourier transform of the equal-time Green's function $G_e(x,0^+) = G_b(x,0^+)$:

$$n_k \propto \int_{-\infty}^{\infty} \frac{e^{ikx}\ dx}{[1-i\Lambda_c x]^2[1+i\Lambda_n x]} \propto -ke^{k/\Lambda_c}\theta(-k) + \frac{\Lambda_n\Lambda_c}{\Lambda_n+\Lambda_c}e^{-k/\Lambda_n}\theta(k). \tag{13}$$

Due to the presence of counter-propagating neutral mode, we have an extra occupation on the $k>0$ side that decays exponentially over the distance $k \sim \Lambda_n$. We believe the discontinuity at $k=0$ is not physical and is an artifact of taking the integration over infinite range. We can instead work with the Green's function for finite circumference L:

$$\begin{aligned} G_\phi(z) &= -2\ln\left(\frac{1-\exp[-2\pi(1-iz\Lambda_c)/\Lambda_c L]}{1-\exp[-2\pi/\Lambda_c L]}\right), \\ G_\eta(w) &= -\ln\left(\frac{1-\exp[-2\pi(1+iw\Lambda_n)/\Lambda_n L]}{1-\exp[-2\pi/\Lambda_n L]}\right). \end{aligned} \tag{14}$$

The corresponding occupation number n_m is given by

$$n_m \propto \int_0^1 \frac{\exp[2\pi imx]\ dx}{[1-\exp(-2\pi(1-i\kappa_c x)/\kappa_c)]^2[1-\exp(-2\pi(1+i\kappa_n x)/\kappa_n)]}. \tag{15}$$

We have introduced $\kappa_{c,n} = \Lambda_{c,n}L$ and measured x in units of L. Expanding the denominators in powers series, one obtains a simple expression for n_m:

$$n_m \propto \sum_{s=0}^{\infty}\sum_{t=0}^{\infty} \delta_{t-s,m}(s+1)e^{-2\pi s/\kappa_c - 2\pi t/\kappa_n}. \tag{16}$$

Normalizing by the value at $m=0$,

$$n_m/n_0 = \begin{cases} e^{-2\pi m/\kappa_n}, & m>0 \\ e^{2\pi m/\kappa_c}\{1-m(1-e^{-2\pi(\kappa_c^{-1}+\kappa_n^{-1})})\}, & m<0. \end{cases} \tag{17}$$

It is exponentially decaying in the $m>0$ region (tail) and linear, of the form $A+Bm$, for $-\kappa_c \ll m < 0$. The coefficients A and B depend on the exact value of the cutoff parameters. To connect with the thermodynamic limit, one can rewrite the occupation numbers in terms of k. We assume that the momentum cut-offs Λ_n and Λ_c remain finite in the $L \to \infty$ limit.

$$n_k/n_0 = \begin{cases} e^{-k/\Lambda_n}, & k>0 \\ e^{k/\Lambda_c}(1-k(\Lambda_c^{-1}+\Lambda_n^{-1})), & k<0. \end{cases} \tag{18}$$

Figure 1 shows occupation numbers for $N=11$ electrons at half-filling obtained from finite-size exact diagonalization[5]. Our trial wave function indicates that the tail region, $x>0$ with $x = kR_{ed}$, does not disappear in the thermodynamic limit. Here $R_{ed} = 2\sqrt{N}$ is the radius of a uniform, compact half-filled droplet. The trial wave function was constructed by occupying about half of the composite fermions in the lowest Landau level, while each of the other half singly occupies the higher Landau levels after the projection. In the Laughlin state all of the composite fermions occupy the lowest Landau level to form an incompressible liquid[15]. Such differences in the occupation behavior of composite fermions underlie the observed difference of electron occupation for compressible/incompressible states. From Eqs. (17) and (18) we expect $n_k = A+Bk$ for small negative k, and we find that this expression (open squares) fits the numerical results well for suitable choice of A and B.

In this paper we have demonstrated that the edge occupation numbers calculated from our trial wave function are similar to those of Lee and Wen's theory. It would be interesting to see if similar excitation properties can be found.

The work of SREY has been supported by the Ministry of Education under Grant No. BSRI-96-2444.

---

[1] B. I. Halperin, P. A. Lee, and N. Read, Phys. Rev. B **47**, 7312 (1993).
[2] N. Read, Semicond. Sci. Technol. **9**, 1859 (1994).
[3] R. Shankar and Ganpathy Murthy, Phys. Rev. Lett. **79**, 4437 (1997); B. I. Halperin and Ady Stern, cond-mat/9802061.
[4] V. Pasquier and F. D. M. Haldane, cond-mat/9712169.
[5] S.-R. Eric Yang and J. H. Han, Phys. Rev. B **57**, R12681 (1998).
[6] D. H. Lee, Phys. Rev. Lett. **80**, 4745 (1998).
[7] N. Read, cond-mat/9804294.
[8] A. V. Shytov, L. S. Levitov, and B. I. Halperin, Phys. Rev. Lett. **80**, 141 (1998).
[9] D. H. Lee and X. G. Wen; cond-mat/9809160.
[10] J. H. Han and S.-R. Eric Yang, Phys. Rev. B **58**, R10163 (1998).
[11] Sami Mitra and A. H. MacDonald, Phys. Rev. **48**, 2005 (1993); E. H. Rezayi and F. D. M. Haldane, Phys. Rev. B **50**, 17199 (1993); S.-R. Eric Yang, Sami Mitra, A. H. MacDonald, and M. P. A. Fisher, J. Korean Phys. Soc. **49**, S10 (1996).
[12] X. G. Wen, Int. J. Mod. Phys. B **6**, 1711 (1992).
[13] M. Grayson, D. C. Tsui, L. N. Pfeiffer, K. W. West, and A. M. Chang, Phys. Rev. Lett. **80**, 1062 (1998).
[14] J. H. Han and D. J. Thouless, Phys. Rev. B **55**, 1926 (1997).
[15] J. K. Jain and R. K. Kamilla, Int. J. Mod. Phys. B **11**, 2621 (1997).



FIG. 1. Occupation numbers for $N = 11(\bullet)$ computed numerically. Occupation numbers for small negative values of x are fitted with a linear function (open squares).